

DOMAIN	QUALITY MEASURE	Full Name	DOMAIN	QUALITY MEASURE	Full Name	
PREVENTIVE AND WELL	W30	Well-Child Visits in the First 30 Months of Life - 6 or more well visits for children 15 months old and 2 or more well visits for children 15-30 months	ACUTE AND CHRONIC CONDITION MANAGEMENT	CWP	Appropriate Testing for Pharyngitis (3 years and older diagnosed with pharyngitis, dispense and antibiotic and received a group A strep test)	
	WCV	Well-Child Visits annually for 3-21 years of age		MMA 50%	Medication Management for People with Asthma- controller medication 50% of treatment period	
	AAP	Adults' Access to Preventive/Ambulatory Health Services (minimum of one visit per calendar year)		MMA 75%	Medication Management for People with Asthma- controller medication 75% of treatment period	
	AWV	Adult Wellness Visit annually		AMR	Asthma Medication Ratio (persistent asthma patients 5-64 years of age and had minimum ratio of .50 controller medications to total asthma medications)	
	HRA- Annually	Health Risk Assessment		PCE	Pharmacotherapy Management of COPD Exacerbation (patients 40 years and older who had an acute IP discharge for ED visit on or between 1/1-11/30 and were dispensed appropriate medications)	
	CIS- MMR	Childhood Immunization Status- Measles, Mumps, Rubella		URI-Peds	Appropriate Treatment for Upper Respiratory Infection (patients 3 months of age and older with diagnosis of URI that did not receive an antibiotic)	
	CIS-VZV	Childhood Immunization Status- Varicella Zoster		ART	Disease Modifying Anti-Rheumatic Drug Therapy for Rheumatoid Arthritis (patients 18 and older diagnosed with RA and who were dispensed at least 1 DMARD in the measurement year)	
	CIS-Comb 10	Childhood Immunization Status- Combination 10 (DTaP, IPV, MMR, Hib, VZV, PCV, HepA, RV, Influenza)		OMW	Osteoporosis Management (women 67-85 who suffered a fracture and had either a BMD test or prescription to treat osteoporosis within 6 months post fracture)	
	LSC	Lead Screening in Children (1 or more lead blood tests prior to 24 months)		LBP	Use of Imaging Studies for Low Back Pain (patients 18-50 with primary diagnosis of low back pain who did not receive imaging within 28 days of the diagnosis)	
	CHL - Adult (21-24)	Chlamydia Screening in Women ages 21-24 (minimum one test annually)		MRP - IP	Medication Reconciliation Within 30 Days Post Inpatient Discharge	
	CHL- Peds (16-20)	Chlamydia Screening in Women ages 16-20 (minimum one test annually)		AAB- acute bronchitis-adult	Avoidance of Antibiotic Treatment for Acute Bronchitis (patients 3 months and older diagnosed with acute bronchitis or bronchiolitis who did not receive an antibiotic)	
	COL	Colorectal Cancer Screening (FOBT annually, FIT-DNA test q3 years, colonoscopy q10 years, flexible sigmoidoscopy or CT colonography q5 years)		CDC (all components)	Comprehensive Diabetes Care (diabetic patients 18-75 years who had annually each of the 6 components listed below)	
	CCS	Cervical Cancer Screening (cervical cytology within last 3 years for women 21-64 or hrHPV testing within last 5 years for women 30-64 or hrHPV co-testing within the last 5 years for women 30-64)		1. DM- Retinopathy/ Eye Exam	Annual Screening for Diabetic Retinal Disease	
	BCS	Breast Cancer Screening for Women 50-74 who have had screening for breast cancer		2. DM- Nephropathy Screening	Annual Nephropathy Screening or Evidence of Nephropathy	
	PPC-Timeliness	Timeliness of prenatal Care (deliveries that received a prenatal care visit in the first trimester)		3. DM- HbA1c	Annual HbA1c test	
	PPC- Postpartum	Postpartum Care (women that had a postpartum visit on or between 7-84 days post delivery)		4. DM- HbA1c Poor Control >9%	Most recent HbA1c test during the measurement year greater than 9% or missing a result	
	FLU	Flu Vaccinations Annually (various age ranges)		5. DM-HbA1c Controlled <8%	Most recent HbA1c test during the measurement year less than 8%	
	TOB	Tobacco Use Screening annually and Cessation Intervention if positive		6. CBP - Diabetic	Most recent blood pressure test during measurement year less than 140/90	
	DSF (CMS Prev 12)	Annual Depression Screening ages 12 years and older and, if positive, follow up within 30 days		KED (For Commercial and Medicaid lives only)	% of members 18-85 with diabetes who received a kidney health evaluation, defined by an estimated glomerular filtration rate (eGFR) and a urine albumin-creatinine ratio (uACR), during measurement year	
	COA- (4 components of measure: Functional Status Assessment, Advanced Care Planning, Medication Review, Pain Assessment)	Care for Older Adults 66 years and older who had each component annually		CBP	Controlling High Blood Pressure (patients 18-85 years with a diagnosed with hypertension and most recent test during measurement year less than 140/90)	
	ABA	Adult Annual BMI with follow-up if outside of parameters		MAH - 2+ fills for 80% PY	Medication Adherence- Hypertension (ages 40-75 years)	
	BMI with f/u	BMI outside of normal parameters have follow up		MAC - 2+ fills for 80% PY	Medication Adherence-Cholesterol	
	FRM	Falls Risk Management annual screening		MAD	Medication Adherence- Diabetes (ages 40-75 years where member remained on statin medication for at least 80% of the treatment period)	
	UTILIZATION	AMB-ER		Ambulatory ER utilization rate per 1000	SPD	Statin Therapy for Patients with Diabetes (ages 40-75 years who were dispensed at least one statin medication during the measurement year)
		ACSC - IP		Ambulatory Care Sensitive Conditions- Inpatient admissions	SPC	Statin Therapy for Patients with Cardiovascular Disease (males 21-75 years or females 40-75 years, diagnosed with ASCVD dispensed at least one high to moderate intensity statin medication during the measurement year)
IP readmit Rate - 30 day			ADD-Initiation	Follow up Care for Children Prescribed ADHD Medication Initiation Phase (ages 6-12 with at least 1 visit in initial 30 days)		
PAER- Adult		Potentially Avoidable Emergency Room Visit- 19 years+	ADD-Continuation	Follow up Care for Children Prescribed ADHD Medication Continuation Phase (ages 6-12 with at least 2 visits in 9 months after Initiation Phase)		
PAER- <18		Potentially Avoidable Emergency Room Visit- 18 years and below	MH-1 (CMS)	Mental Health 1 (patients 18 years and older diagnosed with depression who achieved remission in 12 months (+/- 60 days) after an index event) (Remission is achieved when a PHQ9 score is less than 5)		
Brand Formulary - Rate Adult			SMC Forms	Suspected Medical Condition- Medical Risk Adjustment/HCC		
Brand Formulary- Rate <18			PAF	Patient Assessment Form		
			RISK ADJUSTMENT REQUIREMENT			