

DOMAIN	QUALITY MEASURE	Full Name	DOMAIN	QUALITY MEASURE	Full Name
PREVENTIVE AND WELL	W30	Well-Child Visits in the First 30 Months of Life - 6 or more well visits for children 15 months old and 2 or more well visits for children 15-30 months	ACUTE AND CHRONIC CONDITION MANAGEMENT	CWP	Appropriate Testing for Pharyngitis (3 years and older diagnosed with pharyngitis, dispense and antibiotic and received a group A strep test age and had minimum ratio of .50 controller medications to total asthma medications)
	WCV	Well-Child Visits annually for 3-21 years of age		AMR	
	AAP	Adults' Access to Preventive/Ambulatory Health Services (minimum of one visit per calendar year)		PCE	Pharmacotherapy Management of COPD Exacerbation (patients 40 years and older who had an acute IP discharge for ED visit on or between 1/1-11/30 and were dispensed appropriate medications)
	AWV	Adult Wellness Visit annually		URI-Peds	Appropriate Treatment for Upper Respiratory Infection (patients 3 months of age and older with diagnosis of URI that did not receive an antibiotic)
	CIS- MMR	Childhood Immunization Status- Measles, Mumps, Rubella		ART	Disease Modifying Anti-Rheumatic Drug Therapy for Rheumatoid Arthritis (patients 18 and older diagnosed with RA and who were dispensed at least 1 DMARD in the measurement year)
	CIS-VZV	Childhood Immunization Status- Varicella Zoster		OMW	Osteoporosis Management (women 67-85 who suffered a fracture and had either a BMD test or prescription to treat osteoporosis within 6 months post fracture)
	CIS-Comb 10	Childhood Immunization Status- Combination 10 (DTaP, IPV, MMR, HiB, VZV, PCV, HepA, RV, Influenza)		LBP	Use of Imaging Studies for Low Back Pain (patients 18-74 with primary diagnosis of low back pain who did not receive imaging within 28 days of the diagnosis)
	LSC	Lead Screening in Children (1 or more lead blood tests prior to 24 months)		TRC (Medicare Only)	Transitions of Care - Medicare members 18 and older with notification of IP admission, Receipt of DC information (documentation of receipt of DC info at DC), Patient Engagement post DC w/I 30 days, Med Rec 30 days post DC
	CHL - Adult (21-24)	Chlamydia Screening in Women ages 21-24 (minimum one test annually)		AAB- acute bronchitis-adult	Avoidance of Antibiotic Treatment for Acute Bronchitis (patients 3 months and older diagnosed with acute bronchitis or bronchiolitis who did not receive an antibiotic)
	CHL- Peds (16-20)	Chlamydia Screening in Women ages 16-20 (minimum one test annually)		HBD	Hemoglobin A1c Control - % of members 18-75 w/diabetes whose HbA1c was: * Controlled <8% * Poor controlled >9%
	COL	Colorectal Cancer Screening (FOBT annually, FIT-DNA test q3 years, colonoscopy q10 years, flexible sigmoidoscopy or CT colonography q5 years)		BPD	Blood Pressure Control - % of members 18-75 w/diabetes whose blood pressure was adequately controlled (<140/90 mm Hg) during measurement year
	CCS	Cervical Cancer Screening (cervical cytology within last 3 years for women 21-64 or hrHPV testing within last 5 years for women 30-64 or hrHPV co-testing within the last 5 years for women 30-64)		EED	Eye Exam - % of members 18-75 w/diabetes who had a retinal eye exam
	BCS	Breast Cancer Screening for Women 50-74 who have had screening for breast cancer		KED	Kidney Health Evaluation - % of members 18-85 w/diabetes who received a kidney health evaluation, defined by an estimated glomerular filtration rate (eGFR) and a urine albumin-creatinine ratio (uACR), during measurement year
	PPC-Timeliness	Timeliness of prenatal Care (deliveries that received a prenatal care visit in the first trimester)		CBP	Controlling High Blood Pressure (patients 18-85 years with a diagnosed with hypertension and most recent test during measurement year less than 140/90)
	PPC- Postpartum	Postpartum Care (women that had a postpartum visit on or between 7-84 days post delivery)		MAH - 2+ fills for 80% PY	Medication Adherence- Hypertension (ages 40-75 years)
	FLU	Flu Vaccinations Annually (various age ranges)		MAC - 2+ fills for 80% PY	Medication Adherence-Cholesterol
	TOB	Tobacco Use Screening annually and Cessation Intervention if positive		MAD	Medication Adherence- Diabetes (ages 40-75 years where member remained on statin medication for at least 80% of the treatment period)
	DSF (CMS Prev 12)	Annual Depression Screening ages 12 years and older and, if positive, follow up within 30 days		SPD	Statin Therapy for Patients with Diabetes (ages 40-75 years who were dispensed at least one statin medication during the measurement year)
	COA- (3 components of measure: Functional Status Assessment, Medication Review, Pain Assessment)	Care for Older Adults 66 years and older who had each component annually		SPC	Statin Therapy for Patients with Cardiovascular Disease (males 21-75 years or females 40-75 years, diagnosed with ASCVD dispensed at least one high to moderate intensity statin medication during the measurement year)
	ACP	Advanced Care Planning - % of members 66-80 with advanced illness, an indication of frailty or who are receiving palliative care, and adults 81 or older who had advanced care planning during the measurement year		ADD-Initiation	Follow up Care for Children Prescribed ADHD Medication Initiation Phase (ages 6-12 with at least 1 visit in initial 30 days)
FRM	Falls Risk Management annual screening	ADD-Continuation	Follow up Care for Children Prescribed ADHD Medication Continuation Phase (ages 6-12 with at least 2 visits in 9 months after Initiation Phase)		
UTILIZATION	AMB-ER	Ambulatory ER utilization rate per 1000	RISK ADJUSTMENT REQUIREMENT	MH-1 (CMS)	Mental Health 1 (patients 18 years and older diagnosed with depression who achieved remission in 12 months (+/- 60 days) after an index event) (Remission is achieved when a PHQ9 score is less than 5)
	ACSC - IP	Ambulatory Care Sensitive Conditions- Inpatient admissions		SMC Forms	Suspected Medical Condition- Medical Risk Adjustment/HCC
	IP readmit Rate - 30 day			PAF	Patient Assessment Form
	PAER- Adult	Potentially Avoidable Emergency Room Visit- 19 years+	PLAN REQUIREMENT	HRA- Annually	Health Risk Assessment - School City of Mishawaka
	PAER- <18	Potentially Avoidable Emergency Room Visit- 18 years and below			
	Brand Formulary - Rate Adult				
	Brand Formulary- Rate <18				
	DBO	Deprescribing of Benzodiazepines in Older Adults - % of members 67 or older who were dispensed benzodiazepines and achieved a 20% decrease or greater in benzodiazepine (DME) dose during the measurement year			
AXR	Antibiotic Utilization for Respiratory Conditions - % of episodes for members 3 months or older with diagnosis of respiratory condition that results in an antibiotic dispensing event on or 3 days after Episode Date				